

# INTERNATIONAL SEARCH REPORT

International Application No  
NL/R02004/000012

**A. CLASSIFICATION OF SUBJECT MATTER**  
**IPC 7 G01N33/15 A61K49/00**

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
**IPC 7 G01N A61K**

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE**

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SOMLYAI G ET AL: "THE BIOLOGICAL EFFECTS OF DEUTERIUM-DEPLETED WATER, A POSSIBLE NEW TOOL IN CANCER THERAPY" DEUTSCHE ZEITSCHRIFT FUER ONKOLOGIE, HEIDELBERG, DE, vol. 30, no. 4, 1998, pages 91-94, XP009006973 ISSN: 0931-0037 cited in the application page 92, left-hand column, paragraph 5 - page 93, left-hand column, paragraph 4	1,2
X	WO 95/18545 A (HYD KUTATO-FEJLESZTO KTF; SOMLYAI, GABOR) 13 July 1995 (1995-07-13) cited in the application page 3, line 30 - page 4, line 18	1,2 -/-

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

**17 February 2005**

Date of mailing of the international search report

**28/02/2005**

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

**Albayrak, T**

## INTERNATIONAL SEARCH REPORT

PCT/R02004/000012

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 855 921 A (SOMLYAI ET AL) 5 January 1999 (1999-01-05) column 4, line 54 - column 6, line 63 -----	1,2

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International Application No

PCT/R02004/000012

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 9518545	A	13-07-1995	WO	9518545 A1		13-07-1995
			AT	205055 T		15-09-2001
			AU	691483 B2		21-05-1998
			AU	5977794 A		01-08-1995
			DE	69428213 D1		11-10-2001
			DE	69428213 T2		13-06-2002
			DK	738114 T3		26-11-2001
			EP	0738114 A1		23-10-1996
			FI	962747 A		02-08-1996
			JP	9511129 T		11-11-1997
			RU	2125817 C1		10-02-1999
			SK	87996 A3		05-02-1997
US 5855921	A	05-01-1999	HU	208084 B		30-08-1993
			AT	402692 B		25-07-1997
			AT	902992 A		15-12-1996
			BE	1006186 A3		07-06-1994
			CA	2122612 A1		13-05-1993
			CH	685282 A5		31-05-1995
			CN	1071836 A ,C		12-05-1993
			CZ	9401020 A3		15-12-1994
			DE	4232465 A1		06-05-1993
			DK	170832 B1		05-02-1996
			ES	2077530 A1		16-11-1995
			FI	941979 A		29-06-1994
			FR	2683148 A1		07-05-1993
			GB	2276086 A ,B		21-09-1994
			WO	9308794 A1		13-05-1993
			IT	1255390 B		31-10-1995
			JP	8501275 T		13-02-1996
			JP	3569916 B2		29-09-2004
			KR	253874 B1		15-04-2000
			LU	88175 A1		15-03-1993
			NL	9220026 A ,B,		01-09-1994
			NO	941590 A		23-06-1994
			SE	513624 C2		09-10-2000
			SE	9401406 A		07-06-1994

**AMENDED CLAIMS**

[Received by the International Bureau on 26 April 2005 (26.04.05):  
original claims 1-2 replaced by amended claims 1-2 (2 pages)]

1. The method for in vivo determination on tested animals of the efficient concentration of Deuterium Depleted Water for cancer therapy is characterized by the fact that it provides Deuterium Depleted Water administering to tested animals before and after tumor grafting with animal grafts and it takes the following steps:

- A) Deuterium Depleted Water administering to Wistar outbred rats by diet, with concentration less than 100 ppm, over a period of 60 days, simultaneously to dieting a control group of animals with water having 150 ppm content of Deuterium (tap water), over the same period of time..
- B) Viability determination for the tumor cells to be grafted, using tripan blue
- C) Grafting of the animals in the experimental group and the control animals in the 60th day, subcutaneous, with  $1 \times 10^7$  malign tumor cells in 0,5 ml normal saline solution of 256 Walker sarcoma (the solid tumor) and T8 Guérin lymphotropic epithelioma (solid tumor), both of them having cells with a viability over 98%.
- D) Continuously and long-term administering, by diet, of Deuterium Depleted Water, with concentration less than 100 ppm deuterium, period over which the followings are to be done:
  - a. Starting with the 4-th post-graft day the tumor nodules measurement and examination is performed on each 2-3 days;
  - b. Monitoring of animals physiological condition by weekly weighing, monitoring their food and water consumption, notifying the toxic condition occurrence
  - c. After 60 days, when all the animals in control group are dead, preferable between the 160th and 200th day after graft, the effect produced by administering of established concentration of Deuterium Depleted Water is observed on the surviving animals homeostasis from experimental groups, respectively the way how humoral immune system and cellular immune system of these animals has been influenced, by performing of a series of examination on immunological condition of the animals, namely: leucocytes formula test to establish lymphocytes and blastic cells levels; hematopoietic marrow tests to establish the plasmocytes and NK-K cells levels.
- E) Determination of efficient concentration of Deuterium Depleted Water for tested surviving animals depending on new homeostasis occurrence, and on the results obtained related to tumoral regression, as well as to cancer curing.

2. Method, as per claim no. 1, characterized by the fact that it determines the 60 ppm Deuterium Depleted Water as the concentration that is the most efficient for cancer therapy and prophylaxis by continuously and long-term administering of this type of water as a daily diet.

Technical issue the invention is solving is the establishing of a method for experimental determination in vivo of an efficient Deuterium content in water, in order to obtain optimum results in cancer therapy on rats.

According to the invention, the method consist in Deuterium Depleted Water administering before and after tumor grafting on animals, following the stages below:

- A) Deuterium Depleted Water administering to Wistar outbred rats by diet, with concentration less than 100 ppm, over a period of 60 days, simultaneously to dieting a control group of animals with water having 150 ppm content of Deuterium (tap water), over the same period of time
- B) Viability determination for the tumor cells to be grafted, using tripan blue
- C) Grafting of the animals in the experimental group and the control animals in the 60th day, subcutaneous, with  $1 \times 10^7$  malign tumor cells in 0,5 ml normal saline solution of 256 Walker sarcoma (the solid tumor) and T8 Guérin lymphotropic epithelioma (solid tumor), both of them having cells with a viability over 98%.
- D) Continuously and long-term administering, by diet, of Deuterium Depleted Water, with concentration less than 100 ppm deuterium, period over which the followings are to be done:
  - a) Starting with the 4-th post-graft day the tumor nodules measurement and examination is performed on each 2-3 days;
  - b) Monitoring of animals physiological condition by weekly weighing, monitoring their food and water consumption, notifying the toxic condition occurrence
  - c) After 60 days, when all the animals in control group are dead, preferable between the 160th and 200th day after graft, the effect produced by administering of established concentration of Deuterium Depleted Water is observed on the surviving animals homeostasis from experimental groups, respectively the way how humoral immune system and cellular immune system of these animals has been influenced, by performing of a series of examination on immunological condition of the animals, namely: leucocytes formula test to establish lymphocytes and blastic cells levels; hematopoietic marrow tests to establish the plasmocytes and NK-K cells levels.
- E) Determination of efficient concentration of Deuterium Depleted Water for tested surviving animals depending on new homeostasis occurrence, and on the results obtained related to tumoral regression, as well as to cancer curing.